

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.

**THIS PAGE BLANK (USPTO)**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


(51) International Patent Classification <sup>7</sup> : <b>A61K 35/78, A61P 29/02</b>		A1	(11) International Publication Number: <b>WO 00/56348</b>
		(43) International Publication Date: 28 September 2000 (28.09.00)	
(21) International Application Number: <b>PCT/US00/07085</b>		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 20 March 2000 (20.03.00)		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(30) Priority Data: 09/272,363 19 March 1999 (19.03.99) US			
(71) Applicant (for all designated States except US): MONSANTO COMPANY [US/US]; 800 N. Lindbergh Boulevard, St. Louis, MO 63167 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): CORLEY, David, G. [-/US]; 164 S. Elm Avenue, St. Louis, MO 63119 (US). OBUKOWICZ, Mark [-/US]; 655 N. Kirkwood Road, Kirkwood, MO 63122 (US).			
(74) Agents: PERRY, Lawrence, S. et al.; Fitzpatrick, Cella, Harper & Scinto, 30 Rockefeller Plaza, New York, NY 10112-3801 (US).			

(54) Title: INFLAMMATORY MEDIATION OBTAINED FROM ATRACTYLODES LANCEA

## (57) Abstract

The present invention provides a method for inhibiting the activity of cyclooxygenase-2 and other proinflammatory factors in a mammal. The method comprises administering to the mammal a therapeutically effective or prophylactically effective amount of an organic extract of *Atractylodes lancea*. The inhibitory effect of the organic extract of this invention on cyclooxygenase-2 activity is substantially greater than the inhibitory effect of the organic extract on cyclooxygenase-1 activity. The present invention also

provided a method for treating a mammal having, or at risk for developing, a condition which is benefited by the inhibition of cyclooxygenase-2 or other proinflammatory factors. The method comprises administering to the mammal a therapeutically effective or prophylactically effective amount of the organic extract of *Atractylodes lancea*.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## TITLE

INFLAMMATORY MEDIATION OBTAINED  
FROM ATRACTYLODES LANCEA

5

## Field of the Invention

10 This invention is in the field of nutritional and  
pharmaceutical agents for the management of  
inflammation related conditions. More specifically,  
this invention relates to the use of organic plant  
extracts to inhibit the activity of proinflammatory  
factors in inflamed tissue, as well as to prevent  
15 the incitation of proinflammatory factors in non-  
inflamed tissue.

## Background of the Invention

20 Prostaglandins are known to play an important role  
in several conditions associated with the  
inflammation process. Consequently, significant  
efforts have been directed towards identification of  
agents which are capable of inhibiting prostaglandin  
synthesis.

25

- 2 -

Nonsteroidal antiinflammatory drugs (NSAIDs) are used to reduce pain and swelling associated with the inflammation process. These compounds function by inhibiting the synthesis of prostaglandins.

5 However, because prostaglandins are also involved in maintaining proper gastrointestinal functioning, these compounds can have serious gastrointestinal side effects. Corticosteroids provide an alternative to NSAIDs; however, corticosteroids can  
10 result in even greater side effects than NSAIDs, especially when long term therapy is required.

NSAIDs are cyclooxygenase inhibitors. They interfere with the activity of cyclooxygenase  
15 enzymes and thereby inhibit the synthesis of prostaglandins. It is now known that there are two cyclooxygenase enzymes that are involved in prostaglandin synthesis. The cyclooxygenase enzyme responsible for prostaglandin synthesis in  
20 gastrointestinal tissue is called cyclooxygenase-1 (COX-1), while the cyclooxygenase enzyme responsible for prostaglandin synthesis in inflamed tissue is called cyclooxygenase-2 (COX-2).

25 It has been reported that NSAIDs may selectively inhibit either COX-1 or COX-2 activity. O'Neill et al., *Molec. Pharmacol.*, 45, 245-254 (1994). This suggests that it should be possible to inhibit the activity of COX-2 without significantly inhibiting  
30 COX-1 activity. Thus, it should be possible to manage patients suffering from inflammation related conditions without causing significant gastrointestinal side effects.

35 Organic extracts of *Atractylodes lancea* have been shown to inhibit cyclooxygenase-1 activity in an enzymatic assay. Resch, et al., J. Nat. Prod., 61,

- 3 -

347-350 (1998). Methanol, dichloromethane and n-hexane extracts from rhizomes of *Atractylodes lancea* all showed inhibitory effects on COX-1 activity.

5 The n-hexane extract was reported to contain an atractylochromene with significant COX-1 inhibitory effects. The reference does not disclose or suggest the use of organic extracts of *Atractylodes lancea* as selective COX-2 inhibitors, or administration of any purified and isolated compounds to mammals such as humans.

10 It would be very desirable to provide an agent that selectively inhibits the activity of COX-2 and other proinflammatory factors without significantly affecting the activity of COX-1.

#### SUMMARY OF THE INVENTION

20 The present invention provides a method for inhibiting the activity of cyclooxygenase-2 and other proinflammatory factors in a mammal. The method comprises administering to the mammal a therapeutically-effective or prophylactically-effective amount of an organic extract of

25 *Atractylodes lancea*. The inhibitory effect of the organic extract of this invention on the activity of cyclooxygenase-2 and other proinflammatory factors is substantially greater than the inhibitory effect of the organic extract on cyclooxygenase-1 activity.

30 The present invention also provides a method for managing a condition in a mammal which is benefited by the inhibition of cyclooxygenase-2 or other proinflammatory factors. The method comprises

35 administering to the mammal a therapeutically-effective or prophylactically-effective amount of the organic extract of *Atractylodes lancea*.

- 4 -

## DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "purified" includes partially purified and completely purified. Thus, a  
5 "purified compound" may be either partially purified or completely purified. The term "proinflammatory factor" refers to substances (e.g., cytokines and eicosinoids) and processes (e.g., binding of  
10 inflammatory cells to protein matrixes) that are involved in the inflammation process. As used herein, the term "extract" includes crude extract, purified extract, and purified compounds obtained by purification of the extract.

15 It has been discovered that organic extracts of the rhizomes of the plant *Atractylodes lancea* exhibit selective inhibition of cyclooxygenase-2 (COX-2). The inhibitory effect is selective in that the inhibition of COX-2 is significantly greater than  
20 the inhibition of cyclooxygenase-1 (COX-1). These extracts also have been shown to inhibit other proinflammatory factors.

Consequently, organic extracts of *Atractylodes*  
25 *lancea* rhizomes may be used to selectively inhibit the activity of COX-2 and other proinflammatory factors in a mammal without causing an equivalent inhibition of COX-1 activity. Proinflammatory factors whose activity may be inhibited by the  
30 organic extracts of *Atractylodes lancea* include COX-2 activity, 15-lipoxygenase activity, thromboxane synthetase activity, inflammatory cell adhesion to fibronectin, inflammatory cell adhesion to VCAM-1, IL-1 $\beta$  cytokine release, IL-2 cytokine  
35 release, IL-6 cytokine release, Interferon- $\gamma$  cytokine release, TNF- $\alpha$  cytokine release, TNF- $\alpha$  mediated PGE<sub>2</sub> release, IL-1 $\alpha$  mediated PGE<sub>2</sub> release,



- 5 -

NF-AT transcription of proinflammatory genes, and  
NF- $\kappa$ B transcription of proinflammatory genes.

5 The extracts of this invention may be used to manage  
a mammal having, or at risk for developing, a  
condition which is benefited by the inhibition of  
COX-2 or other proinflammatory factors. Conditions  
which may be benefited by the inhibition of COX-2  
10 and other proinflammatory factors include, for  
example, general inflammation, arthritis, pain and  
cancer.

15 Preferably, the inhibitory effect of the extract of  
*Atractylodes lancea* on COX-2 is at least about two  
times greater than its inhibitory effect on COX-1.  
More preferably, the inhibitory effect on COX-2 is  
at least about 10 times greater than the inhibitory  
effect on COX-1.

20 Those of ordinary skill in the art of preparing  
pharmaceutical formulations can readily formulate  
pharmaceutical compositions having *Atractylodes*  
*lancea* extracts using known excipients (e.g.,  
saline, glucose, starch, etc.). Similarly, those of  
25 ordinary skill in the art of preparing nutritional  
formulations can readily formulate nutritional  
compositions having *Atractylodes lancea* extracts.  
And those of ordinary skill in the art of preparing  
food or food ingredient formulations can readily  
30 formulate food compositions or food ingredient  
compositions having *Atractylodes lancea* extracts.

35 In addition, those of ordinary skill in the art can  
readily determine appropriate dosages that are  
necessary to achieve the desired therapeutic or  
prophylactic effect upon oral, parenteral, rectal  
and other administration forms. Typically, in-vivo

- 6 -

models (i.e., laboratory mammals) are used to determine the appropriate plasma concentrations necessary to achieve a desired mitigation of inflammation related conditions.

5

The present invention provides a method for managing a condition in a mammal which is benefited by the inhibition of cyclooxygenase-2 or other proinflammatory factors. The method comprises administering to the mammal a therapeutically effective or prophylactically effective amount of the organic extract of *Atractylodes lancea*.

10

The organic extracts of the present invention may be obtained by extraction from *Atractylodes lancea*, in particular, from the rhizomes of *Atractylodes lancea*. In a desirable process, rhizomes of *Atractylodes lancea* are ground into a fine powder, the resultant powder is extracted with a solvent, and the extraction solvent is removed from the extract. If desired, the resultant extract may be further purified to yield a purified extract or one or more purified compounds.

15

20

The grinding step may be accomplished by any commonly known method for grinding a plant substance. For example, the rhizomes may be passed through a grinder to obtain a fine powder. After the rhizomes of *Atractylodes lancea* have been ground into a fine powder, they are combined with an extraction solvent.

25

30

The solution is stirred at a temperature, and for a period of time, that is effective to obtain an extract with the desired inhibitory effects on the activity of COX-2 and/or other proinflammatory factors. The solution should not be overheated, as

35

- 7 -

5 this may result in degradation of the extract. The solution may be stirred at a temperature between about room temperature (25°C) and the boiling point of the extraction solvent. Preferably, the solution is stirred at about room temperature.

10 The length of time during which the rhizome powder is exposed to the extraction solvent is not critical. Up to a point, the longer the rhizome powder is exposed to the extraction solvent, the greater is the amount of extract that may be recovered. Preferably, the solution is stirred for at least 1 minute, more preferably for at least 15 minutes, and most preferably for at least 60 minutes.

20 The extraction process of the present invention is desirably carried out using an organic solvent or a mixture of organic solvents. Organic solvents which may be used in the extraction process of the present invention include hydrocarbon solvents, ether solvents, chlorinated solvents, acetone, ethyl acetate, butanol, ethanol, methanol, isopropyl alcohol and mixtures thereof. Hydrocarbon solvents which may be used in the present invention include heptane, hexane and pentane. Ether solvents which may be used in the present invention include diethyl ether. Chlorinated solvents which may be used in the present invention include dichloromethane and chloroform. Preferably, the solvent is a nonpolar organic solvent, such as dichloromethane or hexane.

35 The relative amount of solvent used in the extraction process may vary considerably, depending upon the particular solvent employed. Typically, for each 100 grams of rhizome powder to be

- 8 -

extracted, about 500 ml of extraction solvent would be used.

5 The organic solvent may be removed from the extract by any method well known in the field of chemistry for removing organic solvents from a desired product, including, for example, rotary evaporation.

10 It is believed that the inhibitory effect of the *Atractylodes lancea* extract of this invention on the activity of COX-2 and other proinflammatory factors is due to one or more compounds present in the extract. Compounds present in the extract which  
15 inhibit the activity of COX-2 and other proinflammatory factors may be isolated and purified by those of ordinary skill in the art using methods known in the art. For example, column chromatography and fractional distillation may be  
20 used to obtain pure compounds from the *Atractylodes lancea* extract of this invention.

The isolation and purification of particular compounds from organic extracts of *Atractylodes lancea* may be performed as described in Resch, et  
25 al., J. Nat. Prod., 61, 347-350 (1998), the entire contents of which are incorporated by reference herein. The methods disclosed therein may be used to isolate and purify compounds which exhibit selective inhibitory activity of COX-2 and other  
30 proinflammatory factors.

The examples which follow are intended to illustrate certain preferred embodiments of the invention, and no limitation of the invention is implied.

35

- 9 -

## PREPARATION EXAMPLE 1

5 Rhizomes of *Atractylodes lancea* (Oriental Ginseng and Gift, St. Louis, MO) were dried and sliced. The sliced rhizomes were ground into a fine powder using a coffee grinder. 100 grams of the resulting powder were added to 500 ml of dichloromethane and stirred at room temperature for 1 hour. The solvent was then removed by rotary evaporation, leaving 5.1  
10 grams of a yellow-brown oil as the extract.

## EXAMPLE 1

15 Inhibitory Effect of Organic Extract of *Atractylodes lancea* Rhizomes on COX-1 and COX-2 Activities

The organic extract obtained in Preparation Example 1 was evaluated for selective inhibition of COX-1 and COX-2. The COX-1 and COX-2 inhibition  
20 activities were determined in vitro by the art-recognized method described by Gierse et al., *J. Biochem.*, 305, 479-484 (1995), summarized below.

25 Preparation of recombinant COX baculoviruses

Recombinant COX-1 was prepared by cloning a 2.0 kb fragment containing the coding region of human or murine COX-1 into a BamH1 site of the baculovirus transfer vector pVL1393 (Invitrogen) to generate the  
30 baculovirus transfer vectors for COX-1 in a manner similar to the method of D.R. O'Reilly et al., *Baculovirus Expression Vectors: A Laboratory Manual* (1992).

35 Recombinant baculoviruses were isolated by transfecting 4  $\mu$ g of baculovirus transfer vector DNA into ( $2 \times 10^8$ ) SF9 insect cells along with 200 mg of

- 10 -

linearized baculovirus plasmid DNA by the calcium phosphate method. (See M.D. Summers and G.E. Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agric. Exp. Station Bull. 1555 (1987)). Recombinant viruses were purified by three rounds of plaque purification and high titer ( $10^7$ - $10^8$  pfu/ml) stocks of virus were prepared.

For large scale production, SF9 insect cells were infected in 10 liter fermentors ( $0.5 \times 10^6$ /ml) with the recombinant baculovirus stock such that the multiplicity of infection was 0.1. After 72 hours the cells were centrifuged and the cell pellet was homogenized in Tris/Sucrose (50 mM: 25%, pH 8.0) containing 1% 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS). The homogenate was centrifuged at 10,000 X G for 30 minutes, and the resultant supernatant was stored at  $-80^\circ\text{C}$ .

Recombinant COX-2 was prepared by cloning a 2.0kb fragment containing the coding region of human or murine COX-2 in the same manner as described above.

#### Assay for COX-1 and COX-2 Activities

COX-1 and COX-2 activities were assayed as  $\text{PGE}_2$  formed/ $\mu\text{g}$  protein/time using ELISA to detect prostaglandin  $\text{E}_2$  synthesized from arachidonic acid. CHAPS-solubilized insect cell membranes containing the COX-1 or COX-2 enzyme were incubated in a potassium phosphate buffer (50 mM, pH 8.0) containing epinephrine, phenol, and heme. Compounds were pre-incubated with the appropriate enzyme for 10-20 minutes. Arachidonic acid ( $10 \mu\text{M}$ ) was then

- 11 -

added to the mixture and the reaction was permitted to occur for ten minutes at room temperature (25°C).

Any reaction between the arachidonic acid and the enzyme was stopped after ten minutes by transferring 40  $\mu$ l of reaction mixture into 160  $\mu$ l ELISA buffer and 25  $\mu$ M indomethacin. The PGE<sub>2</sub> formed was measured by standard ELISA technology (Cayman Chemical).

A 200 mg sample of the extract obtained from Preparation Example 1 was dissolved in 2 ml of dimethyl sulfoxide (DMSO) for bioassay testing to determine the COX-1 and COX-2 inhibitory effects of the extract. The results of these bioassays are reported in Table 1.

Table 1

Amount of Extract (mg/ml)	Percent Relative Control (COX-1)	Percent Relative Control (COX-2)
0.50	33.3	4.9
0.17	86.3	10.9
0.056	155.2	17.8
0.019	152.2	25.4
0.0062	125.5	47.3
0.0021	104.7	63.8
0.00069	83.6	52.4
0.000023	77.2	79.8
0.000076	72.6	66.4
0.000025	66.0	61.4

Figure 1 is graph showing the data of Table 1. As can be seen in Figure 1, the extract obtained in Example 1 has a much greater inhibitory effect on COX-2 than it does on COX-1. Figure 1 shows that the

- 12 -

COX-1  $IC_{50}$  of the extract was 350  $\mu\text{g/ml}$ , whereas the  
COX-2  $IC_{50}$  of the extract was 5  $\mu\text{g/ml}$ . Thus, the  
*Atractylodes lancea* extract of Example 1 had an  
inhibitory effect on COX-2 that was 70 times greater  
than its inhibitory effect on COX-1.

A sample of the organic extract of Preparation  
Example 1 in isopropanol (1 mg/ml) was prepared for  
HPLC analysis. The results of the HPLC analysis of  
Preparation Example 1 are shown in Figure 2.

#### PREPARATION EXAMPLE 2

Rhizomes of *Atractylodes lancea* (East Earth Herb,  
Eugene, OR) were dried and sliced. The sliced  
rhizomes were ground into a fine powder using a  
coffee grinder. 100 grams of the resulting powder  
were added to 500 ml of dichloromethane and stirred  
at room temperature for 1 hour. The solvent was  
then removed by rotary evaporation, leaving 5.1  
grams of a yellow-brown oil.

#### EXAMPLE 2

A 200 mg sample of the extract obtained from  
Preparation Example 2 was dissolved in 2 ml of  
dimethyl sulfoxide (DMSO) and subjected to bioassay  
testing in the same manner as employed in Example 1.  
The results of these bioassays are reported in  
Table 2.



- 13 -

Table 2

5	Amount of Extract (mg/ml)	Percent Relative Control (COX-1)	Percent Relative Control (COX-2)
	0.50	16.3	4.1
	0.17	49.4	11.7
	0.056	75.7	21.6
	0.019	85.9	33.5
10	0.0062	84.7	45.5
	0.0021	84.3	54.0
	0.00069	73.1	59.1
	0.00023	69.0	77.9
	0.000076	64.3	52.9
15	0.000025	63.8	78.2

Figure 3 is a graph showing the data of Table 2. As can be seen in Figure 3, the extract obtained in Example 2 has a much greater inhibitory effect on COX-2 than it does on COX-1. Figure 3 shows that the COX-1  $IC_{50}$  of the extract was 150  $\mu$ g/ml, whereas the COX-2  $IC_{50}$  of the extract was 4  $\mu$ g/ml. Thus, the *Atractylodes lancea* extract of Example 2 had an inhibitory effect on COX-2 that was 37 times greater than its inhibitory effect on COX-1.

A sample of the organic extract of Preparation Example 2 in isopropanol (1 mg/ml) was prepared for HPLC analysis. The results of the HPLC analysis of Preparation Example 2 are shown in Figure 4.

### EXAMPLE 3

A 200 mg sample of the *Atractylodes lancea* extract obtained from Preparation Example 2 was dissolved in 2 ml of DMSO. A 100  $\mu$ l sample of the resultant

- 14 -

solution was fractionated over a 500 mg C-18 Bond Elute SPE column as follows: the column was equilibrated with 100% methanol, followed by a 1:1 mixture of water to methanol. The 100  $\mu$ l DMSO extract solution was added to 2.0 ml of a 1:1 water to methanol solution, and the resultant solution was added to the column. The column was then eluted with an additional 1.0 ml of a 1:1 mixture of water to methanol, thereby yielding a first fraction eluted with 3.0 ml of 1:1 water to methanol. The next fraction was eluted with 3.0 ml of a 1:4 water to methanol solution. The third fraction was eluted with 3.0 ml of pure methanol and the last fraction was eluted with 3.0 ml of dichloromethane. The four fractions were dried by rotary evaporation, and then the four fractions along with a crude *Atractylodes lancea* extract sample were dissolved in 100  $\mu$ l DMSO for COX-2 bioassays. The results of these bioassays are reported in Table 3.

Table 3

Elution Solvent	COX-2 IC <sub>50</sub> (100 $\mu$ g/ml)
1:1 H <sub>2</sub> O:CH <sub>3</sub> OH	> 200 $\mu$ g/ml
1:4 H <sub>2</sub> O:CH <sub>3</sub> OH	> 200 $\mu$ g/ml
100% CH <sub>3</sub> OH	18 $\mu$ g/ml
100% CH <sub>2</sub> Cl <sub>2</sub>	> 200 $\mu$ g/ml
Crude extract	16 $\mu$ g/ml

Figure 5 is a graph showing the data of Table 3. As can be seen in Figure 5, the pure methanol fraction displayed greater COX-2 inhibitory effects than any of the other purified fractions. The crude extract also displayed significant COX-2 inhibitory effects.

- 15 -

The pure methanol fraction and the positive control (i.e., crude extract) were analyzed using HPLC. The results of the HPLC analyses of the methanol fraction and the positive control are shown in Figures 6 and 7, respectively.

## EXAMPLE 4

A 5g sample of the *Atractylodes lancea* extract of Preparation Example 1 was dissolved in 50 ml of dichloromethane and chromatographed over 100g SiO<sub>2</sub> vacuum flash. After the 50 ml extract solution was applied to the column, the column was eluted with the following 200 ml fractions: 100% dichloromethane; 9:1 dichloromethane to methanol; 7:3 dichloromethane to methanol; 3:7 dichloromethane to methanol and 100% methanol. The five fractions were dried by rotary evaporation, and then the five fractions and a positive control (i.e., crude extract) were dissolved in DMSO (100 µg/ml) for COX-2 bioassays. The results of the bioassays are reported in Table 4.

Table 4

Elution Solvent	COX-2 IC <sub>50</sub> (100 µg/ml) in DMSO
100% CH <sub>2</sub> Cl <sub>2</sub>	> 500 µg/ml
9:1 CH <sub>2</sub> Cl <sub>2</sub> :CH <sub>3</sub> OH	50 µg/ml
7:3 CH <sub>2</sub> Cl <sub>2</sub> :CH <sub>3</sub> OH	6 µg/ml
3:7 CH <sub>2</sub> Cl <sub>2</sub> :CH <sub>3</sub> OH	5 µg/ml
100% CH <sub>3</sub> OH	50 µg/ml
Crude extract	9 µg/ml

- 16 -

Figure 8 is a graph showing the data of Table 4. As can be seen in Figure 8, the third and fourth fractions (7:3 CH<sub>2</sub>Cl<sub>2</sub> to CH<sub>3</sub>OH and 3:7 CH<sub>2</sub>Cl<sub>2</sub> to CH<sub>3</sub>OH, respectively) displayed significant COX-2 inhibitory activity.

The 7:3 CH<sub>2</sub>Cl<sub>2</sub> to CH<sub>3</sub>OH fraction was analyzed by HPLC. The results of the HPLC analysis are shown in Figure 9.

#### EXAMPLE 5

##### Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on 15-Lipoxygenase Activity

The inhibitory effect of the organic extract obtained in Preparation Example 1 on 15-lipoxygenase was evaluated using methods described in Auerback et al., Anal. Biochem., 201, 375-380 (1992). A test solution of the extract was prepared at 100 µg/ml in 0.1% DMSO.

A 100 µg/ml sample of the test solution was incubated with 15 U of 15-lipoxygenase (obtained from rabbit reticulocytes) in a phosphate buffered saline solution at a pH of 7.4 and a temperature of 4°C. The reaction was initiated by addition of 256 µM linoleic acid as substrate and run for 10 minutes after which the reaction was terminated by addition of N-benzoyl leucomethylene blue (LMB). The level of 15-HETE was determined by measuring absorbance at 660 nm. The results of these assays are set forth in Table 5.

- 17 -

## EXAMPLE 6

Inhibitory Effect Of Organic Extract of *Atractylodes*  
*lancea* Rhizomes on Thromboxane Synthetase Activity

5

The inhibitory effect of the organic extract obtained in Preparation Example 1 on Thromboxane Synthetase was evaluated using methods described in Fiddler et al., Circulation, 81 (Suppl) I69-I78  
10 (1990). A test solution of the extract was prepared at 100  $\mu\text{g/ml}$  in 0.1% DMSO.

A 100  $\mu\text{g/ml}$  sample of the test solution was incubated with 1:200 dilution of thromboxane  $A_2$   
15 synthase (isolated from a microsomal fraction of rabbit platelets) and 5 ng prostaglandin  $G_2$  as substrate in Tris buffer at a pH of 7.5 for 30 minutes at a temperature of 37°C. The thromboxane  $A_2$  formed was immediately converted to thromboxane  $B_2$   
20 which was quantitated by a radioimmunoassay. The results of these assays are set forth in Table 5.

## EXAMPLE 7

25 Inhibitory Effect Of Organic Extract of *Atractylodes*  
*lancea* Rhizomes on Fibronectin Mediated Cell  
Adhesion

The inhibitory effect of the organic extract obtained in Preparation Example 1 on Fibronectin  
30 mediated cell adhesion was evaluated using methods described in Nowlin, et al., J. Biol. Chem., 268, 20352-20359 (1993). Test solutions of the extract were prepared at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  
35  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$  in 0.1% DMSO.

- 18 -

These assays measured the adhesion of NRK 2 (normal rat kidney) cells to a fibronectin-coated well. Each of the test solutions in modified MEM-HEPES buffer at a pH of 7.4 was incubated for 30 minutes at a temperature of 37°C. The reactions were initiated by addition of NRK 2 cells ( $2 \times 10^6/\text{ml}$ ) and incubated for 30 minutes. Each well was then washed 6 times with Dulbecco's PBS followed by addition of  $5\mu\text{M}$  calcein AM and a further 2 hour incubation period. Quantitation of fluorescent intensity resulting from interaction of calcein AM with cells attached to the fibronectin coated plate was read with a Cytofluor 2300 plate reader with B filter excitation at 485 nm and emission at 530 nm. The results of these assays are set forth in Table 5.

## EXAMPLE 8

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on VCAM-1 Mediated Cell Adhesion

The inhibitory effect of the organic extract obtained in Preparation Example 1 on VCAM-1 mediated cell adhesion was evaluated using methods described in Stoltenborg, et al., J. Immunological Methods, 175, 59-68 (1994). Test solutions of the extract were prepared at  $100\mu\text{g}/\text{ml}$ ,  $10\mu\text{g}/\text{ml}$ ,  $1\mu\text{g}/\text{ml}$ ,  $0.1\mu\text{g}/\text{ml}$  and  $0.01\mu\text{g}/\text{ml}$  in 0.1% DMSO.

These assays measured adherence of Jurkat (human T lymphoid) cells to wells coated with recombinant human VCAM-1 isolated from the membranes of infected SF9 cells. On the day of the assay, the plates were washed with Dubelco's PBS (DPBS) at a pH of 7.2. Each of the test solutions was incubated with Jurkat cells ( $0.5$  to  $1 \times 10^6/\text{ml}$  labeled with  $5\mu\text{g}/\text{ml}$  calcein AM) and  $2.5\text{ ng}/\text{ml}$  PMA (phorbol 12-myristate 13-

- 19 -

acetate) in RPMI at a pH of 7.5 in the coated wells. The plates were incubated at 25°C for 60 minutes without shaking and washed with DPBS. Adhesion was then quantitated by reading plates on a Cytoflour 2300. The results of these assays are set forth in Table 5.

## EXAMPLE 9

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on IL-1 $\beta$  Cytokine Release

The inhibitory effect of the organic extract obtained in Preparation Example 1 on IL-1 $\beta$  cytokine release was evaluated using methods described in Welker, et al., International Arch of Allergy and Immunology, 109, 110-115 (1996). Test solutions of the extract were prepared at 100  $\mu$ g/ml, 10  $\mu$ g/ml, 1  $\mu$ g/ml, 0.1  $\mu$ g/ml and 0.01  $\mu$ g/ml in 0.1% DMSO.

Each of the test solutions was incubated overnight with 25 ng/ml lipopolysaccharide (LPS) and stimulated human peripheral blood mononuclear leukocytes (PBMNL) in growth medium RPMI-1640 at a pH of 7.4 and a temperature of 37°C. The IL-1 $\beta$  cytokine production levels in the conditioned medium were quantitated using a sandwich ELISA kit. The results of these assays are set forth in Table 5.

EXAMPLE 10

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on IL-2 Cytokine Release

The inhibitory effect of the organic extract obtained in Preparation Example 1 on IL-2 cytokine release was evaluated using methods described in

- 20 -

Welker, et al., International Arch of Allergy and Immunology, 109, 110-115 (1996). Test solutions of the extract were prepared at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$  in 0.1% DMSO.

5

Each of the test solutions was incubated with 10  $\mu\text{g/ml}$  Concanavalin-A (Con-A) and stimulated human peripheral blood mononuclear leukocytes (PBMNL) in growth medium RPMI-1640 at a pH of 7.4 and a temperature of 37°C. The IL-2 cytokine production levels in the conditioned medium were quantitated using a sandwich ELISA kit. The results of these assays are set forth in Table 5.

10

15

## EXAMPLE 11

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on IL-2 Cytokine Release

20

The inhibitory effect of the organic extract obtained in Preparation Example 1 on IL-2 cytokine release also was evaluated using methods described in Koizumi, et al., 103, 469-475 (1986), which employ trypsinized Jurkat cells rather than PBMNL. Test solutions of the extract were prepared at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$  in 0.1% DMSO.

25

30

35

Each of the test solutions was incubated overnight, in the presence or absence of co-stimulation by 1  $\mu\text{g/ml}$  calcium ionophore (A23187) and 25 ng/ml PMA (phorbol 12-myristate 13-acetate), with trypsinized Jurkat (human T lymphoid) cells ( $2 \times 10^6/\text{ml}$ ) suspended with 10% fetal bovine serum in RPMI-1640 at a pH of 7.4 and a temperature of 37°C. The cell suspension was subjected to centrifugation, and the supernatant was evaluated for IL-2 release by use of



- 21 -

an IL-2 immunoassay kit. The results of these assays are set forth in Table 5.

## EXAMPLE 12

5

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on IL-6 Cytokine Release

10 The inhibitory effect of the organic extract obtained in Preparation Example 1 on IL-6 cytokine release was evaluated using methods described in Welker, et al., International Arch of Allergy and Immunology, 109, 110-115 (1996). Test solutions of the extract were prepared at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ ,  
15 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$  in 0.1% DMSO.

Each of the test solutions was incubated overnight with LPS (25 ng/ml) and stimulated human peripheral blood mononuclear leukocytes (PBMNL) in growth  
20 medium RPMI-1640 at a pH of 7.4 and a temperature of 37°C. The IL-6 cytokine production levels in the conditioned medium were quantitated using a sandwich ELISA kit. The results of these assays are set forth in Table 5.

25

## EXAMPLE 13

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on Interferon- $\gamma$  Cytokine Release

30

The inhibitory effect of the organic extract obtained in Preparation Example 1 on Interferon- $\gamma$  (IFN- $\gamma$ ) cytokine release was evaluated using methods described in the following references: (1) Cohen,  
35 et al., Am. J. Clin. Pathol. 105, 589-598 (1996); (2) Henderson, et al., TIPS, 13, 145-151 (1992); (3) Welker, et al., International Arch of Allergy and

- 22 -

Immunology, 109, 110-115 (1996); and (4) Elias, et al., J. Immunol., 138, 3812-3816 (1987).

Test solutions of the extract were prepared at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$  in 0.1% DMSO.

Each of the test solutions was incubated overnight with Concanavalin A (Con-A, 10  $\mu\text{g/ml}$ )-stimulated human peripheral blood mononuclear cells (PBMNCs) in RPMI-1640 growth medium at a pH of 7.4 and a temperature of 37°C. IFN- $\gamma$  cytokine levels in the conditioned medium were then quantitated using a sandwich ELISA kit. The results of these assays are set forth in Table 5.

#### EXAMPLE 14

##### Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on TNF- $\alpha$ Cytokine Release

The inhibitory effect of the organic extract obtained in Preparation Example 1 on TNF- $\alpha$  cytokine release was evaluated using methods described in the following references: (1) Cohen, et al., Am. J. Clin. Pathol. 105, 589-598 (1996); (2) Henderson, et al., TIPS, 13, 145-151 (1992); and (3) Welker, et al., International Arch of Allergy and Immunology, 109, 110-115 (1996). Test solutions of the extract were prepared at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$  in 0.1% DMSO.

Each of the test solutions was incubated overnight with LPS (25 ng/ml) and stimulated human peripheral blood mononuclear cells (PBMNCs) in RPMI-1640 growth medium at a pH of 7.4 and a temperature of 37°C. TNF- $\alpha$  cytokine levels in the conditioned medium were

- 23 -

then quantitated using a sandwich ELISA kit. The results of these assays are set forth in Table 5.

## EXAMPLE 15

5

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on TNF- $\alpha$  Mediated PGE<sub>2</sub> Release

10 The inhibitory effect of the organic extract obtained in Preparation Example 1 on TNF- $\alpha$  mediated PGE<sub>2</sub> release was evaluated using methods described in Lenardo, et al., Cell, 58, 227-229 (1989). Test solutions of the extract were prepared at 100  $\mu$ g/ml, 10  $\mu$ g/ml, 1  $\mu$ g/ml, 0.1  $\mu$ g/ml and 0.01  $\mu$ g/ml  
15 in 0.1% DMSO.

Each of the test solutions was incubated overnight in the presence or absence of 25 nM Tumor Necrosis Factor- $\alpha$  (TNF- $\alpha$ ), with trypsinized HeLa (human  
20 epithelioid cervix carcinoma) S3 cells ( $2 \times 10^6$ /ml) suspended with 10% fetal bovine serum in MEM at a pH of 7.3 and a temperature of 37°C. The cell suspension of each well was then transferred to an Eppendorf vial, subjected to centrifugation, and the  
25 supernatant was evaluated for released PGE<sub>2</sub> (prostaglandin E<sub>2</sub>) by radioimmunoassay. The results of these assays are set forth in Table 5.

## EXAMPLE 16

30

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on IL-1 $\alpha$  Mediated PGE<sub>2</sub> Release

35 The inhibitory effect of the organic extract obtained in Preparation Example 1 on IL-1 $\alpha$  mediated PGE<sub>2</sub> release was evaluated using methods described in

- 24 -

Maloff, et al., Clin. Chim. Acta, 180 73-78 (1989).  
Test solutions of the extract were prepared at 100  
 $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$   
in 0.1% DMSO.

5 Each of the test solutions was incubated overnight,  
in the presence or absence of 1 nM interleukin- $1\alpha$   
(IL- $1\alpha$ ), with trypsinized WI-38 (human diploid  
fibroblast lung) cells ( $10^6/\text{ml}$ ) suspended with 10%  
10 fetal bovine serum in MEM at a pH of 7.3 and a  
temperature of 37°C. The cell suspension of each  
well was then transferred to an Eppendorf vial,  
subjected to centrifugation, and the supernatant was  
evaluated for released PGE $_2$  (prostaglandin E $_2$ ) by  
15 radioimmunoassay. The results of these assays are  
set forth in Table 5.

## EXAMPLE 17

20 Inhibitory Effect Of Organic Extract of *Atractylodes*  
*lancea* Rhizomes on NF $\kappa$ B

The inhibitory effect of the organic extract  
obtained in Preparation Example 1 on NF $\kappa$ B was  
25 evaluated using the methods described in Karttunen,  
et al., Proc. Nat'l. Acad. Sci. USA, 88, 3972-3976  
(1991). Test solutions of the extract were prepared  
at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  
 $\mu\text{g/ml}$  in 0.1% DMSO.

30 Jurkat (human T lymphoid) cells transfected with a  
response element-lacZ reporter in which  
transcription of the  $\beta$ -galactosidase gene is  
directed by the binding site for the NF- $\kappa$ B  
35 transcription factor ( $\kappa$ B-Z cells) were used in these  
assays. Each of the test solutions was incubated  
with  $\kappa$ B-Z cells ( $2 \times 10^5$ ) in the presence of 2  $\mu\text{M}$  of

- 25 -

a calcium ionophore (A23187) and 20 ng/ml PMA in RPMI buffer at a pH of 7.4 for 4 hours at 37°C. The cells were then centrifuged, resuspended in buffer and conversion of FDG (fluorescein di- $\beta$ -D-galactopyranoside) to fluorescein by induced  $\beta$ -galactosidase activity was determined after overnight incubation in the dark at 25°C. Fluorescence intensity was measured using Cytofluor (2300) plate reader with excitation at 485 nm and emission at 530 nm. The results of these assays are set forth in Table 5.

## EXAMPLE 18

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on NF-AT

The inhibitory effect of the organic extract obtained in Preparation Example 1 on NF-AT was evaluated using the methods described in Emmel, et al., Science, 246, 1617-1620 (1989). Test solutions of the extract were prepared at 100  $\mu$ g/ml, 10  $\mu$ g/ml, 1  $\mu$ g/ml, 0.1  $\mu$ g/ml and 0.01  $\mu$ g/ml in 0.1% DMSO.

Jurkat (human T lymphoid) cells transfected with a response element-lacZ reporter in which transcription of the  $\beta$ -galactosidase gene is directed by the binding site for the NFAT-1 transcription factor were used for the assays. Each of the test solutions was incubated with cells ( $2 \times 10^5$ ) in the presence of 2  $\mu$ M of a calcium ionophore (A23187) and 20 ng/ml PMA in RPMI buffer at a pH of 7.4 for 4 hours at 37°C. The cells were then centrifuged and resuspended in buffer, and conversion of FDG (fluorescein di- $\beta$ -D-galactopyranoside) to fluorescein by induced  $\beta$ -galactosidase activity was determined after

- 26 -

overnight incubation in the dark at 25°C.  
 Fluorescence intensity was measured using a  
 Cytofluor (2300) plate reader with excitation at 485  
 nm and emission at 530 nm. The results of these  
 assays are set forth in Table 5.

Table 5

Proinflammatory Factor	A. lancea extract conc. (µg/ml)	% Inhibition	IC <sub>50</sub> (µg/ml)
15-Lipoxygenase	100	50	
Thromboxane Synthetase	100	88	
Cell Adhesion (Fibronectin)	100	116	5.4
Cell Adhesion (Fibronectin)	10.0	65	
Cell Adhesion (Fibronectin)	1.0	13	
Cell Adhesion (Fibronectin)	0.1	15	
Cell Adhesion (Fibronectin)	0.01	10	
Cell Adhesion (VCAM-1)	100	109	9.9
Cell Adhesion (VCAM-1)	10.0	52	
Cell Adhesion (VCAM-1)	1.0	-11	
Cell Adhesion (VCAM-1)	0.1	9	
Cell Adhesion (VCAM-1)	0.01	13	
Cytokine Release (IL-1β)	100	110	7.2
Cytokine Release (IL-1β)	10.0	85	
Cytokine Release (IL-1β)	1.0	-8	
Cytokine Release (IL-1β)	0.1	-3	
Cytokine Release (IL-1β)	0.01	-9	
Cytokine Release (IL-2) PBMNL	100	102	0.4549
Cytokine Release (IL-2) PBMNL	10.0	89	
Cytokine Release (IL-2) PBMNL	1.0	45	
Cytokine Release (IL-2) PBMNL	0.1	37	
Cytokine Release (IL-2) PBMNL	0.01	20	

- 27 -

	Proinflammatory Factor	A. lancea extract conc. ( $\mu\text{g/ml}$ )	% Inhibition	IC <sub>50</sub> ( $\mu\text{g/ml}$ )
5	Cytokine Release (IL-2) Jurkat	100	87	70.6
	Cytokine Release (IL-2) Jurkat	10.0	-13	
	Cytokine Release (IL-2) Jurkat	1.0	-11	
	Cytokine Release (IL-2) Jurkat	0.1	-9	
	Cytokine Release (IL-2) Jurkat	0.01	-9	
10	Cytokine Release (IL-6)	100	100	2.7
	Cytokine Release (IL-6)	10.0	99	
	Cytokine Release (IL-6)	1.0	3	
	Cytokine Release (IL-6)	0.1	-4	
	Cytokine Release (IL-6)	0.01	-2	
15	Cytokine Release (Interferon- $\gamma$ )	100	99	5.7
	Cytokine Release (Interferon- $\gamma$ )	10.0	68	
	Cytokine Release (Interferon- $\gamma$ )	1.0	9	
	Cytokine Release (Interferon- $\gamma$ )	0.1	-2	
	Cytokine Release (Interferon- $\gamma$ )	0.01	4	
20	Cytokine Release (TNF- $\alpha$ )	100	107	9.3
	Cytokine Release (TNF- $\alpha$ )	10.0	59	
	Cytokine Release (TNF- $\alpha$ )	1.0	-18	
	Cytokine Release (TNF- $\alpha$ )	0.1	-13	
	Cytokine Release (TNF- $\alpha$ )	0.01	-2	
25	TNF- $\alpha$ Mediated PGE <sub>2</sub> Release	100	90	
	TNF- $\alpha$ Mediated PGE <sub>2</sub> Release	10.0	20	
	TNF- $\alpha$ Mediated PGE <sub>2</sub> Release	1.0	-5	
	TNF- $\alpha$ Mediated PGE <sub>2</sub> Release	0.1	-2	
	TNF- $\alpha$ Mediated PGE <sub>2</sub> Release	0.01	0	
30	IL-1 $\alpha$ Mediated PGE <sub>2</sub> Release	100	110	
	IL-1 $\alpha$ Mediated PGE <sub>2</sub> Release	10.0	44	
	IL-1 $\alpha$ Mediated PGE <sub>2</sub> Release	1.0	20	
	IL-1 $\alpha$ Mediated PGE <sub>2</sub> Release	0.1	8	
	IL-1 $\alpha$ Mediated PGE <sub>2</sub> Release	0.01	-1	

- 28 -

Proinflammatory Factor	A. lancea extract conc. ( $\mu\text{g/ml}$ )	% Inhibition	IC <sub>50</sub> ( $\mu\text{g/ml}$ )
NF $\kappa$ B transcription	100	96	7
NF $\kappa$ B transcription	10.0	59	
NF $\kappa$ B transcription	1.0	11	
NF $\kappa$ B transcription	0.1	7	
NF $\kappa$ B transcription	0.01	-13	
NF-AT transcription	100	96	8.8
NF-AT transcription	10.0	54	
NF-AT transcription	1.0	7	
NF-AT transcription	0.1	-12	
NF-AT transcription	0.01	-10	

## EXAMPLE 19

A mammal having or at risk for developing a condition which is benefited by inhibition of the activity of COX-2 or other proinflammatory factors may be treated with a therapeutically or prophylactically effective amount of the *Atractylodes lancea* extract of this invention.

The therapeutically effective or prophylactically effective amount of *Atractylodes lancea* extract may be administered to the mammal by any of the known routes for administering a pharmaceutical agent to a patient, including parenterally, orally and rectally. The *Atractylodes lancea* extract may be administered using a dosing regimen which is effective to inhibit the activity of COX-2 or other proinflammatory factors in the mammal for a desired period of time. The proper dosage amount may be determined by routine experimentation using methods known in the art.



- 29 -

Other variations and modifications of this invention will be obvious to those skilled in the art. This invention is not limited, except as set forth in the claims.

- 30 -

## WHAT IS CLAIMED IS:

1. A method for inhibiting the activity of a proinflammatory factor in a mammal, said method comprising the step of administering to the mammal an organic extract of *Atractylodes lancea*.
2. A method according to claim 1, wherein said proinflammatory factor is selected from the group consisting of cyclooxygenase-2, 15-lipoxygenase, thromboxane synthetase, inflammatory cell adhesion to fibronectin, inflammatory cell adhesion to VCAM-1, IL-1 $\beta$  cytokine release, IL-2 cytokine release, IL-6 cytokine release, Interferon- $\gamma$  cytokine release, TNF- $\alpha$  cytokine release, TNF- $\alpha$  mediated PGE<sub>2</sub> release, IL-1 $\alpha$  mediated PGE<sub>2</sub> release, NF-AT transcription of proinflammatory genes, and NF- $\kappa$ B transcription of proinflammatory genes.
3. A method according to claim 1, wherein said proinflammatory factor is cyclooxygenase-2.
4. A method according to claim 3, wherein the inhibitory effect of said extract on cyclooxygenase-2 is greater than or equal to about 2 times greater than the inhibitory effect of said extract on cyclooxygenase-1.
5. A method according to claim 3, wherein the inhibitory effect of said extract on cyclooxygenase-2 is greater than or equal to about 10 times greater than the inhibitory effect of said extract on cyclooxygenase-1.
6. A method according to any of claims 1-5, wherein said organic extract is a purified compound obtained by a process comprising the steps of:

- 31 -

(i) exposing rhizomes of *Atractylodes lancea* to an organic solvent under conditions appropriate to remove a proinflammatory factor inhibitory extract from said rhizomes; and

(ii) isolating said proinflammatory factor inhibitory extract.

7. A method according to claim 6, wherein step (i) comprising mixing said rhizomes with said solvent and stirring the resultant mixture at a temperature between about 25°C and the boiling point of said solvent for at least one minute.

8. A method according to claim 7, wherein said solvent is an organic solvent.

9. A method according to claim 8, wherein said organic solvent is selected from the group consisting of hydrocarbon solvents, ethers, chlorinated solvents, acetone, ethyl acetate, butanol, ethanol, methanol, isopropyl alcohol and mixtures thereof.

10. A method according to claim 9, wherein said nonpolar organic solvent is dichloromethane.

11. A method according to claim 9, wherein step (ii) comprises separating said solvent from said organic extract by evaporating said solvent.

12. A method according to claim 6, comprising administering a therapeutically effective or prophylactically effective amount of said purified compound to a mammal which has or is at risk for developing a condition which is benefited by the inhibition of the activity of a proinflammatory factor.

- 32 -

13. A method according to claim 12, wherein said condition is a cyclooxygenase-2 mediated condition.

14. A method according to claim 13, wherein said cyclooxygenase-2 mediated condition is general inflammation.

15. A method according to claim 13, wherein said cyclooxygenase-2 mediated condition is arthritis.

16. A method according to claim 13, wherein said cyclooxygenase-2 mediated condition is pain.

17. A method according to claim 13, wherein said cyclooxygenase-2 mediated condition is cancer.

18. A method according to claim 12, wherein said purified compound is administered as a pharmaceutical composition comprising a pharmaceutically acceptable excipient.

19. A method according to claim 12, wherein said purified compound is administered as a nutritional composition comprising a nutritionally acceptable excipient.

20. A method according to claim 12, wherein said purified compound is administered as a food composition.

21. A method according to claim 12, wherein said purified compound is administered as a food ingredient composition.

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6







FIGURE 7







# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/US 00/07085

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A61K35/78 A61P29/02

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 613 591 A (ABURADA MASAKI ET AL) 23 September 1986 (1986-09-23) column 2, line 38 - line 50; claims ---	1-21
X	DATABASE WPI Section Ch, Week 199237 Derwent Publications Ltd., London, GB; Class B03, AN 1992-303541 XP002142228 & JP 04 208222 A (TSUMURA & CO), 29 July 1992 (1992-07-29) abstract --- -/--	1-21

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

10 July 2000

Date of mailing of the international search report

25/07/2000

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Seegert, K

# INTERNATIONAL SEARCH REPORT

In ational Application No

PCT/US 00/07085

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI  Section Ch, Week 199208  Derwent Publications Ltd., London, GB;  Class B04, AN 1992-060692  XP002142229  &amp; JP 04 005237 A (NONOKAWA SHOJI YG),  9 January 1992 (1992-01-09)  abstract</p>	1-21
Y	<p>RESCH M. ET AL: "5-Lipoxygenase and  cyclooxygenase-1 inhibitory active  compounds from Atractylodes lancea"  J. NAT. PROD.,  vol. 61, 1998, pages 347-250, XP000916172  cited in the application  abstract</p>	1-21
Y	<p>DATABASE MEDLINE 'Online!  AN: 94178130, 1993  XP002142227  abstract  &amp; CHANG M L ET AL: "The influence of  Chinese traditional medicine on the  production and activity of interleukin 1  (IL-1)"  JOURNAL OF MICROBIOLOGY AND IMMUNOLOGY ,  vol. 26, no. 1, 1993, pages 15-24,</p>	1-21

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/07085

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4613591 A	23-09-1986	JP 1633501 C	20-01-1992
		JP 2056329 B	29-11-1990
		JP 59036619 A	28-02-1984
		AT 383740 B	10-08-1987
		AT 299783 A	15-01-1987
		BE 897403 A	14-11-1983
		CH 659948 A	13-03-1987
		DE 3328263 A	23-02-1984
		FR 2531864 A	24-02-1984
		GB 2126085 A,B	21-03-1984
		HU 187103 B	28-11-1985
		IT 1169771 B	03-06-1987
		NL 8302391 A	16-03-1984
JP 4208222 A	29-07-1992	NONE	
JP 4005237 A	09-01-1992	NONE	

Form PCT/ISA/210 (patent family annex) (July 1992)

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 September 2000 (28.09.2000)

PCT

(10) International Publication Number  
**WO 00/56348 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 35/78**,  
A61P 29/02

G. [—/US]: 164 S. Elm Avenue, St. Louis, MO 63119 (US). **OBUKOWICZ, Mark** [—/US]: 655 N. Kirkwood Road, Kirkwood, MO 63122 (US).

(21) International Application Number: PCT/US00/07085

(22) International Filing Date: 20 March 2000 (20.03.2000)

(74) Agents: **BLOSSER, Harley, G.** et al.: Senniger, Powers, Leavitt & Roedel, One Metropolitan Square, 16th Floor, St. Louis, MO 63102 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/272,363 19 March 1999 (19.03.1999) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): **PHARMACIA CORPORATION** [US/US]: Corporate Patent Department, P.O. Box 5110, Chicago, IL 60680 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW). Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM). European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,


(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **CORLEY, David**,

[Continued on next page]

(54) Title: INFLAMMATORY MEDIATION OBTAINED FROM *ATRACYLODES LANCEA*

A. LANCEA



(57) Abstract: The present invention provides a method for inhibiting the activity of cyclooxygenase-2 and other proinflammatory factors in a mammal. The method comprises administering to the mammal a therapeutically effective or prophylactically effective amount of an organic extract of *Atracylodes lancea*. The inhibitory effect of the organic extract of this invention on cyclooxygenase-2 activity is substantially greater than the inhibitory effect of the organic extract on cyclooxygenase-1 activity. The present invention also provided a method for treating a mammal having, or at risk for developing, a condition which is benefited by the inhibition of cyclooxygenase-2 or other proinflammatory factors. The method comprises administering to the mammal a therapeutically effective or prophylactically effective amount of the organic extract of *Atracylodes lancea*.

WO 00/56348 A1



MC, NL, PT, SE). OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**(15) Information about Correction:**

see PCT Gazette No. 26/2002 of 27 June 2002, Section II

**Published:**

— with international search report

**(48) Date of publication of this corrected version:**

27 June 2002

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## TITLE

INFLAMMATORY MEDIATION OBTAINED  
FROM ATRACTYLODES LANCEA

5

## Field of the Invention

This invention is in the field of nutritional and  
pharmaceutical agents for the management of  
10 inflammation related conditions. More specifically,  
this invention relates to the use of organic plant  
extracts to inhibit the activity of proinflammatory  
factors in inflamed tissue, as well as to prevent  
the incitation of proinflammatory factors in non-  
15 inflamed tissue.

## Background of the Invention

Prostaglandins are known to play an important role  
20 in several conditions associated with the  
inflammation process. Consequently, significant  
efforts have been directed towards identification of  
agents which are capable of inhibiting prostaglandin  
synthesis.

25

- 2 -

Nonsteroidal antiinflammatory drugs (NSAIDs) are used to reduce pain and swelling associated with the inflammation process. These compounds function by inhibiting the synthesis of prostaglandins.

5 However, because prostaglandins are also involved in maintaining proper gastrointestinal functioning, these compounds can have serious gastrointestinal side effects. Corticosteroids provide an alternative to NSAIDs; however, corticosteroids can  
10 result in even greater side effects than NSAIDs, especially when long term therapy is required.

NSAIDs are cyclooxygenase inhibitors. They interfere with the activity of cyclooxygenase  
15 enzymes and thereby inhibit the synthesis of prostaglandins. It is now known that there are two cyclooxygenase enzymes that are involved in prostaglandin synthesis. The cyclooxygenase enzyme responsible for prostaglandin synthesis in  
20 gastrointestinal tissue is called cyclooxygenase-1 (COX-1), while the cyclooxygenase enzyme responsible for prostaglandin synthesis in inflamed tissue is called cyclooxygenase-2 (COX-2).

25 It has been reported that NSAIDs may selectively inhibit either COX-1 or COX-2 activity. O'Neill et al., *Molec. Pharmacol.*, 45, 245-254 (1994). This suggests that it should be possible to inhibit the activity of COX-2 without significantly inhibiting  
30 COX-1 activity. Thus, it should be possible to manage patients suffering from inflammation related conditions without causing significant gastrointestinal side effects.

35 Organic extracts of *Atractylodes lancea* have been shown to inhibit cyclooxygenase-1 activity in an enzymatic assay. Resch, et al., *J. Nat. Prod.*, 61,



- 3 -

347-350 (1998). Methanol, dichloromethane and n-hexane extracts from rhizomes of *Atractylodes lancea* all showed inhibitory effects on COX-1 activity.

5 The n-hexane extract was reported to contain an atractylochromene with significant COX-1 inhibitory effects. The reference does not disclose or suggest the use of organic extracts of *Atractylodes lancea* as selective COX-2 inhibitors, or administration of any purified and isolated compounds to mammals such as humans.

10 It would be very desirable to provide an agent that selectively inhibits the activity of COX-2 and other proinflammatory factors without significantly affecting the activity of COX-1.

#### SUMMARY OF THE INVENTION

20 The present invention provides a method for inhibiting the activity of cyclooxygenase-2 and other proinflammatory factors in a mammal. The method comprises administering to the mammal a therapeutically-effective or prophylactically-effective amount of an organic extract of

25 *Atractylodes lancea*. The inhibitory effect of the organic extract of this invention on the activity of cyclooxygenase-2 and other proinflammatory factors is substantially greater than the inhibitory effect of the organic extract on cyclooxygenase-1 activity.

30 The present invention also provides a method for managing a condition in a mammal which is benefited by the inhibition of cyclooxygenase-2 or other proinflammatory factors. The method comprises

35 administering to the mammal a therapeutically-effective or prophylactically-effective amount of the organic extract of *Atractylodes lancea*.

- 4 -

## DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "purified" includes partially purified and completely purified. Thus, a  
5 "purified compound" may be either partially purified or completely purified. The term "proinflammatory factor" refers to substances (e.g., cytokines and eicosinoids) and processes (e.g., binding of  
10 inflammatory cells to protein matrixes) that are involved in the inflammation process. As used herein, the term "extract" includes crude extract, purified extract, and purified compounds obtained by purification of the extract.

15 It has been discovered that organic extracts of the rhizomes of the plant *Atractylodes lancea* exhibit selective inhibition of cyclooxygenase-2 (COX-2). The inhibitory effect is selective in that the  
20 inhibition of COX-2 is significantly greater than the inhibition of cyclooxygenase-1 (COX-1). These extracts also have been shown to inhibit other proinflammatory factors.

Consequently, organic extracts of *Atractylodes*  
25 *lancea* rhizomes may be used to selectively inhibit the activity of COX-2 and other proinflammatory factors in a mammal without causing an equivalent inhibition of COX-1 activity. Proinflammatory factors whose activity may be inhibited by the  
30 organic extracts of *Atractylodes lancea* include COX-2 activity, 15-lipoxygenase activity, thromboxane synthetase activity, inflammatory cell adhesion to fibronectin, inflammatory cell adhesion to VCAM-1, IL-1 $\beta$  cytokine release, IL-2 cytokine  
35 release, IL-6 cytokine release, Interferon- $\gamma$  cytokine release, TNF- $\alpha$  cytokine release, TNF- $\alpha$  mediated PGE<sub>2</sub> release, IL-1 $\alpha$  mediated PGE<sub>2</sub> release,

- 5 -

NF-AT transcription of proinflammatory genes, and  
NF- $\kappa$ B transcription of proinflammatory genes.

5 The extracts of this invention may be used to manage  
a mammal having, or at risk for developing, a  
condition which is benefited by the inhibition of  
COX-2 or other proinflammatory factors. Conditions  
which may be benefited by the inhibition of COX-2  
and other proinflammatory factors include, for  
10 example, general inflammation, arthritis, pain and  
cancer.

Preferably, the inhibitory effect of the extract of  
*Atractylodes lancea* on COX-2 is at least about two  
15 times greater than its inhibitory effect on COX-1.  
More preferably, the inhibitory effect on COX-2 is  
at least about 10 times greater than the inhibitory  
effect on COX-1.

20 Those of ordinary skill in the art of preparing  
pharmaceutical formulations can readily formulate  
pharmaceutical compositions having *Atractylodes*  
*lancea* extracts using known excipients (e.g.,  
saline, glucose, starch, etc.). Similarly, those of  
25 ordinary skill in the art of preparing nutritional  
formulations can readily formulate nutritional  
compositions having *Atractylodes lancea* extracts.  
And those of ordinary skill in the art of preparing  
food or food ingredient formulations can readily  
30 formulate food compositions or food ingredient  
compositions having *Atractylodes lancea* extracts.

In addition, those of ordinary skill in the art can  
readily determine appropriate dosages that are  
35 necessary to achieve the desired therapeutic or  
prophylactic effect upon oral, parenteral, rectal  
and other administration forms. Typically, in-vivo

- 6 -

models (i.e., laboratory mammals) are used to determine the appropriate plasma concentrations necessary to achieve a desired mitigation of inflammation related conditions.

5

The present invention provides a method for managing a condition in a mammal which is benefited by the inhibition of cyclooxygenase-2 or other proinflammatory factors. The method comprises administering to the mammal a therapeutically effective or prophylactically effective amount of the organic extract of *Atractylodes lancea*.

10

The organic extracts of the present invention may be obtained by extraction from *Atractylodes lancea*, in particular, from the rhizomes of *Atractylodes lancea*. In a desirable process, rhizomes of *Atractylodes lancea* are ground into a fine powder, the resultant powder is extracted with a solvent, and the extraction solvent is removed from the extract. If desired, the resultant extract may be further purified to yield a purified extract or one or more purified compounds.

15

20

25

30

The grinding step may be accomplished by any commonly known method for grinding a plant substance. For example, the rhizomes may be passed through a grinder to obtain a fine powder. After the rhizomes of *Atractylodes lancea* have been ground into a fine powder, they are combined with an extraction solvent.

35

The solution is stirred at a temperature, and for a period of time, that is effective to obtain an extract with the desired inhibitory effects on the activity of COX-2 and/or other proinflammatory factors. The solution should not be overheated, as

- 7 -

this may result in degradation of the extract. The solution may be stirred at a temperature between about room temperature (25°C) and the boiling point of the extraction solvent. Preferably, the solution is stirred at about room temperature.

The length of time during which the rhizome powder is exposed to the extraction solvent is not critical. Up to a point, the longer the rhizome powder is exposed to the extraction solvent, the greater is the amount of extract that may be recovered. Preferably, the solution is stirred for at least 1 minute, more preferably for at least 15 minutes, and most preferably for at least 60 minutes.

The extraction process of the present invention is desirably carried out using an organic solvent or a mixture of organic solvents. Organic solvents which may be used in the extraction process of the present invention include hydrocarbon solvents, ether solvents, chlorinated solvents, acetone, ethyl acetate, butanol, ethanol, methanol, isopropyl alcohol and mixtures thereof. Hydrocarbon solvents which may be used in the present invention include heptane, hexane and pentane. Ether solvents which may be used in the present invention include diethyl ether. Chlorinated solvents which may be used in the present invention include dichloromethane and chloroform. Preferably, the solvent is a nonpolar organic solvent, such as dichloromethane or hexane.

The relative amount of solvent used in the extraction process may vary considerably, depending upon the particular solvent employed. Typically, for each 100 grams of rhizome powder to be

- 8 -

extracted, about 500 ml of extraction solvent would be used.

5 The organic solvent may be removed from the extract by any method well known in the field of chemistry for removing organic solvents from a desired product, including, for example, rotary evaporation.

10 It is believed that the inhibitory effect of the *Atractylodes lancea* extract of this invention on the activity of COX-2 and other proinflammatory factors is due to one or more compounds present in the extract. Compounds present in the extract which  
15 inhibit the activity of COX-2 and other proinflammatory factors may be isolated and purified by those of ordinary skill in the art using methods known in the art. For example, column chromatography and fractional distillation may be  
20 used to obtain pure compounds from the *Atractylodes lancea* extract of this invention.

The isolation and purification of particular compounds from organic extracts of *Atractylodes lancea* may be performed as described in Resch, et  
25 al., J. Nat. Prod., 61, 347-350 (1998), the entire contents of which are incorporated by reference herein. The methods disclosed therein may be used to isolate and purify compounds which exhibit  
30 selective inhibitory activity of COX-2 and other proinflammatory factors.

The examples which follow are intended to illustrate certain preferred embodiments of the invention, and no limitation of the invention is implied.

35

- 9 -

## PREPARATION EXAMPLE 1

5 Rhizomes of *Atractylodes lancea* (Oriental Ginseng and Gift, St. Louis, MO) were dried and sliced. The sliced rhizomes were ground into a fine powder using a coffee grinder. 100 grams of the resulting powder were added to 500 ml of dichloromethane and stirred at room temperature for 1 hour. The solvent was then removed by rotary evaporation, leaving 5.1  
10 grams of a yellow-brown oil as the extract.

## EXAMPLE 1

15 Inhibitory Effect of Organic Extract of *Atractylodes lancea* Rhizomes on COX-1 and COX-2 Activities

The organic extract obtained in Preparation Example 1 was evaluated for selective inhibition of COX-1 and COX-2. The COX-1 and COX-2 inhibition  
20 activities were determined in vitro by the art-recognized method described by Gierse et al., *J. Biochem.*, 305, 479-484 (1995), summarized below.

25 Preparation of recombinant COX baculoviruses

Recombinant COX-1 was prepared by cloning a 2.0 kb fragment containing the coding region of human or murine COX-1 into a BamH1 site of the baculovirus transfer vector pVL1393 (Invitrogen) to generate the  
30 baculovirus transfer vectors for COX-1 in a manner similar to the method of D.R. O'Reilly et al., *Baculovirus Expression Vectors: A Laboratory Manual* (1992).

35 Recombinant baculoviruses were isolated by transfecting 4  $\mu$ g of baculovirus transfer vector DNA into ( $2 \times 10^8$ ) SF9 insect cells along with 200 mg of

- 10 -

linearized baculovirus plasmid DNA by the calcium phosphate method. (See M.D. Summers and G.E. Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agric. Exp. Station Bull. 1555 (1987)). Recombinant viruses were purified by three rounds of plaque purification and high titer ( $10^7$ - $10^8$  pfu/ml) stocks of virus were prepared.

For large scale production, SF9 insect cells were infected in 10 liter fermentors ( $0.5 \times 10^6$ /ml) with the recombinant baculovirus stock such that the multiplicity of infection was 0.1. After 72 hours the cells were centrifuged and the cell pellet was homogenized in Tris/Sucrose (50 mM: 25%, pH 8.0) containing 1% 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS). The homogenate was centrifuged at 10,000 X G for 30 minutes, and the resultant supernatant was stored at  $-80^\circ\text{C}$ .

Recombinant COX-2 was prepared by cloning a 2.0kb fragment containing the coding region of human or murine COX-2 in the same manner as described above.

#### Assay for COX-1 and COX-2 Activities

COX-1 and COX-2 activities were assayed as  $\text{PGE}_2$  formed/ $\mu\text{g}$  protein/time using ELISA to detect prostaglandin  $\text{E}_2$  synthesized from arachidonic acid. CHAPS-solubilized insect cell membranes containing the COX-1 or COX-2 enzyme were incubated in a potassium phosphate buffer (50 mM, pH 8.0) containing epinephrine, phenol, and heme. Compounds were pre-incubated with the appropriate enzyme for 10-20 minutes. Arachidonic acid ( $10 \mu\text{M}$ ) was then



- 11 -

added to the mixture and the reaction was permitted to occur for ten minutes at room temperature (25°C).

Any reaction between the arachidonic acid and the enzyme was stopped after ten minutes by transferring 40  $\mu$ l of reaction mixture into 160  $\mu$ l ELISA buffer and 25  $\mu$ M indomethacin. The PGE<sub>2</sub> formed was measured by standard ELISA technology (Cayman Chemical).

A 200 mg sample of the extract obtained from Preparation Example 1 was dissolved in 2 ml of dimethyl sulfoxide (DMSO) for bioassay testing to determine the COX-1 and COX-2 inhibitory effects of the extract. The results of these bioassays are reported in Table 1.

Table 1

Amount of Extract (mg/ml)	Percent Relative Control (COX-1)	Percent Relative Control (COX-2)
0.50	33.3	4.9
0.17	86.3	10.9
0.056	155.2	17.8
0.019	152.2	25.4
0.0062	125.5	47.3
0.0021	104.7	63.8
0.00069	83.6	52.4
0.000023	77.2	79.8
0.000076	72.6	66.4
0.000025	66.0	61.4

Figure 1 is graph showing the data of Table 1. As can be seen in Figure 1, the extract obtained in Example 1 has a much greater inhibitory effect on COX-2 than it does on COX-1. Figure 1 shows that the

- 12 -

COX-1  $IC_{50}$  of the extract was 350  $\mu$ g/ml, whereas the  
COX-2  $IC_{50}$  of the extract was 5  $\mu$ g/ml. Thus, the  
*Atractylodes lancea* extract of Example 1 had an  
inhibitory effect on COX-2 that was 70 times greater  
than its inhibitory effect on COX-1.

A sample of the organic extract of Preparation  
Example 1 in isopropanol (1 mg/ml) was prepared for  
HPLC analysis. The results of the HPLC analysis of  
Preparation Example 1 are shown in Figure 2.

#### PREPARATION EXAMPLE 2

Rhizomes of *Atractylodes lancea* (East Earth Herb,  
Eugene, OR) were dried and sliced. The sliced  
rhizomes were ground into a fine powder using a  
coffee grinder. 100 grams of the resulting powder  
were added to 500 ml of dichloromethane and stirred  
at room temperature for 1 hour. The solvent was  
then removed by rotary evaporation, leaving 5.1  
grams of a yellow-brown oil.

#### EXAMPLE 2

A 200 mg sample of the extract obtained from  
Preparation Example 2 was dissolved in 2 ml of  
dimethyl sulfoxide (DMSO) and subjected to bioassay  
testing in the same manner as employed in Example 1.  
The results of these bioassays are reported in  
Table 2.

- 13 -

Table 2

Amount of Extract (mg/ml)	Percent Relative Control (COX-1)	Percent Relative Control (COX-2)
0.50	16.3	4.1
0.17	49.4	11.7
0.056	75.7	21.6
0.019	85.9	33.5
0.0062	84.7	45.5
0.0021	84.3	54.0
0.00069	73.1	59.1
0.00023	69.0	77.9
0.000076	64.3	52.9
0.000025	63.8	78.2

Figure 3 is a graph showing the data of Table 2. As can be seen in Figure 3, the extract obtained in Example 2 has a much greater inhibitory effect on COX-2 than it does on COX-1. Figure 3 shows that the COX-1  $IC_{50}$  of the extract was 150  $\mu$ g/ml, whereas the COX-2  $IC_{50}$  of the extract was 4  $\mu$ g/ml. Thus, the *Atractylodes lancea* extract of Example 2 had an inhibitory effect on COX-2 that was 37 times greater than its inhibitory effect on COX-1.

A sample of the organic extract of Preparation Example 2 in isopropanol (1 mg/ml) was prepared for HPLC analysis. The results of the HPLC analysis of Preparation Example 2 are shown in Figure 4.

## EXAMPLE 3

A 200 mg sample of the *Atractylodes lancea* extract obtained from Preparation Example 2 was dissolved in 2 ml of DMSO. A 100  $\mu$ l sample of the resultant

- 14 -

solution was fractionated over a 500 mg C-18 Bond Elute SPE column as follows: the column was equilibrated with 100% methanol, followed by a 1:1 mixture of water to methanol. The 100  $\mu$ l DMSO extract solution was added to 2.0 ml of a 1:1 water to methanol solution, and the resultant solution was added to the column. The column was then eluted with an additional 1.0 ml of a 1:1 mixture of water to methanol, thereby yielding a first fraction eluted with 3.0 ml of 1:1 water to methanol. The next fraction was eluted with 3.0 ml of a 1:4 water to methanol solution. The third fraction was eluted with 3.0 ml of pure methanol and the last fraction was eluted with 3.0 ml of dichloromethane. The four fractions were dried by rotary evaporation, and then the four fractions along with a crude *Atractylodes lancea* extract sample were dissolved in 100  $\mu$ l DMSO for COX-2 bioassays. The results of these bioassays are reported in Table 3.

Table 3

Elution Solvent	COX-2 IC <sub>50</sub> (100 $\mu$ g/ml)
1:1 H <sub>2</sub> O:CH <sub>3</sub> OH	> 200 $\mu$ g/ml
1:4 H <sub>2</sub> O:CH <sub>3</sub> OH	> 200 $\mu$ g/ml
100% CH <sub>3</sub> OH	18 $\mu$ g/ml
100% CH <sub>2</sub> Cl <sub>2</sub>	> 200 $\mu$ g/ml
Crude extract	16 $\mu$ g/ml

Figure 5 is a graph showing the data of Table 3. As can be seen in Figure 5, the pure methanol fraction displayed greater COX-2 inhibitory effects than any of the other purified fractions. The crude extract also displayed significant COX-2 inhibitory effects.

- 15 -

The pure methanol fraction and the positive control (i.e., crude extract) were analyzed using HPLC. The results of the HPLC analyses of the methanol fraction and the positive control are shown in Figures 6 and 7, respectively.

## EXAMPLE 4

A 5g sample of the *Atractylodes lancea* extract of Preparation Example 1 was dissolved in 50 ml of dichloromethane and chromatographed over 100g SiO<sub>2</sub> vacuum flash. After the 50 ml extract solution was applied to the column, the column was eluted with the following 200 ml fractions: 100% dichloromethane; 9:1 dichloromethane to methanol; 7:3 dichloromethane to methanol; 3:7 dichloromethane to methanol and 100% methanol. The five fractions were dried by rotary evaporation, and then the five fractions and a positive control (i.e., crude extract) were dissolved in DMSO (100 µg/ml) for COX-2 bioassays. The results of the bioassays are reported in Table 4.

Table 4

Elution Solvent	COX-2 IC <sub>50</sub> (100 µg/ml) in DMSO
100% CH <sub>2</sub> Cl <sub>2</sub>	> 500 µg/ml
9:1 CH <sub>2</sub> Cl <sub>2</sub> :CH <sub>3</sub> OH	50 µg/ml
7:3 CH <sub>2</sub> Cl <sub>2</sub> :CH <sub>3</sub> OH	6 µg/ml
3:7 CH <sub>2</sub> Cl <sub>2</sub> :CH <sub>3</sub> OH	5 µg/ml
100% CH <sub>3</sub> OH	50 µg/ml
Crude extract	9 µg/ml

- 16 -

Figure 8 is a graph showing the data of Table 4. As can be seen in Figure 8, the third and fourth fractions (7:3 CH<sub>2</sub>Cl<sub>2</sub> to CH<sub>3</sub>OH and 3:7 CH<sub>2</sub>Cl<sub>2</sub> to CH<sub>3</sub>OH, respectively) displayed significant COX-2 inhibitory activity.

The 7:3 CH<sub>2</sub>Cl<sub>2</sub> to CH<sub>3</sub>OH fraction was analyzed by HPLC. The results of the HPLC analysis are shown in Figure 9.

#### EXAMPLE 5

##### Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on 15-Lipoxygenase Activity

The inhibitory effect of the organic extract obtained in Preparation Example 1 on 15-lipoxygenase was evaluated using methods described in Auerback et al., Anal. Biochem., 201, 375-380 (1992). A test solution of the extract was prepared at 100 µg/ml in 0.1% DMSO.

A 100 µg/ml sample of the test solution was incubated with 15 U of 15-lipoxygenase (obtained from rabbit reticulocytes) in a phosphate buffered saline solution at a pH of 7.4 and a temperature of 4°C. The reaction was initiated by addition of 256 µM linoleic acid as substrate and run for 10 minutes after which the reaction was terminated by addition of N-benzoyl leucomethylene blue (LMB). The level of 15-HETE was determined by measuring absorbance at 660 nm. The results of these assays are set forth in Table 5.

- 17 -

## EXAMPLE 6

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on Thromboxane Synthetase Activity

5

The inhibitory effect of the organic extract obtained in Preparation Example 1 on Thromboxane Synthetase was evaluated using methods described in Fiddler et al., Circulation, 81 (Suppl) I69-I78 (1990). A test solution of the extract was prepared at 100  $\mu\text{g/ml}$  in 0.1% DMSO.

10

A 100  $\mu\text{g/ml}$  sample of the test solution was incubated with 1:200 dilution of thromboxane  $\text{A}_2$  synthase (isolated from a microsomal fraction of rabbit platelets) and 5 ng prostaglandin  $\text{G}_2$  as substrate in Tris buffer at a pH of 7.5 for 30 minutes at a temperature of 37°C. The thromboxane  $\text{A}_2$  formed was immediately converted to thromboxane  $\text{B}_2$  which was quantitated by a radioimmunoassay. The results of these assays are set forth in Table 5.

15

20

## EXAMPLE 7

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on Fibronectin Mediated Cell Adhesion

25

The inhibitory effect of the organic extract obtained in Preparation Example 1 on Fibronectin mediated cell adhesion was evaluated using methods described in Nowlin, et al., J. Biol. Chem., 268, 20352-20359 (1993). Test solutions of the extract were prepared at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$  in 0.1% DMSO.

30

35

- 18 -

These assays measured the adhesion of NRK 2 (normal rat kidney) cells to a fibronectin-coated well. Each of the test solutions in modified MEM-HEPES buffer at a pH of 7.4 was incubated for 30 minutes at a temperature of 37°C. The reactions were initiated by addition of NRK 2 cells ( $2 \times 10^6$ /ml) and incubated for 30 minutes. Each well was then washed 6 times with Dulbecco's PBS followed by addition of  $5\mu\text{M}$  calcein AM and a further 2 hour incubation period. Quantitation of fluorescent intensity resulting from interaction of calcein AM with cells attached to the fibronectin coated plate was read with a Cytofluor 2300 plate reader with B filter excitation at 485 nm and emission at 530 nm. The results of these assays are set forth in Table 5.

## EXAMPLE 8

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on VCAM-1 Mediated Cell Adhesion

The inhibitory effect of the organic extract obtained in Preparation Example 1 on VCAM-1 mediated cell adhesion was evaluated using methods described in Stoltenborg, et al., J. Immunological Methods, 175, 59-68 (1994). Test solutions of the extract were prepared at 100  $\mu\text{g}/\text{ml}$ , 10  $\mu\text{g}/\text{ml}$ , 1  $\mu\text{g}/\text{ml}$ , 0.1  $\mu\text{g}/\text{ml}$  and 0.01  $\mu\text{g}/\text{ml}$  in 0.1% DMSO.

These assays measured adherence of Jurkat (human T lymphoid) cells to wells coated with recombinant human VCAM-1 isolated from the membranes of infected SF9 cells. On the day of the assay, the plates were washed with Dubelco's PBS (DPBS) at a pH of 7.2. Each of the test solutions was incubated with Jurkat cells (0.5 to  $1 \times 10^6$ /ml labeled with 5  $\mu\text{g}/\text{ml}$  calcein AM) and 2.5 ng/ml PMA (phorbol 12-myristate 13-



acetate) in RPMI at a pH of 7.5 in the coated wells. The plates were incubated at 25°C for 60 minutes without shaking and washed with DPBS. Adhesion was then quantitated by reading plates on a Cytoflour 2300. The results of these assays are set forth in Table 5.

### EXAMPLE 9

10        Inhibitory Effect Of Organic Extract of *Atractylodes*  
              *lancea* Rhizomes on IL-1 $\beta$  Cytokine Release

15 The inhibitory effect of the organic extract obtained in Preparation Example 1 on IL-1 $\beta$  cytokine release was evaluated using methods described in Welker, et al., International Arch of Allergy and Immunology, 109, 110-115 (1996). Test solutions of the extract were prepared at 100  $\mu$ g/ml, 10  $\mu$ g/ml, 1  $\mu$ g/ml, 0.1  $\mu$ g/ml and 0.01  $\mu$ g/ml in 0.1% DMSO.

Each of the test solutions was incubated overnight with 25 ng/ml lipopolysaccharide (LPS) and stimulated human peripheral blood mononuclear leukocytes (PBMNL) in growth medium RPMI-1640 at a pH of 7.4 and a temperature of 37°C. The IL-1 $\beta$  cytokine production levels in the conditioned medium were quantitated using a sandwich ELISA kit. The results of these assays are set forth in Table 5.

30

EXAMPLE 10

# Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on IL-2 Cytokine Release

35 The inhibitory effect of the organic extract obtained in Preparation Example 1 on IL-2 cytokine release was evaluated using methods described in

- 20 -

Welker, et al., International Arch of Allergy and Immunology, 109, 110-115 (1996). Test solutions of the extract were prepared at 100 µg/ml, 10 µg/ml, 1 µg/ml, 0.1 µg/ml and 0.01 µg/ml in 0.1% DMSO.

5

Each of the test solutions was incubated with 10 µg/ml Concanavalin-A (Con-A) and stimulated human peripheral blood mononuclear leukocytes (PBMNL) in growth medium RPMI-1640 at a pH of 7.4 and a temperature of 37°C. The IL-2 cytokine production levels in the conditioned medium were quantitated using a sandwich ELISA kit. The results of these assays are set forth in Table 5.

10

15

## EXAMPLE 11

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on IL-2 Cytokine Release

20

25

The inhibitory effect of the organic extract obtained in Preparation Example 1 on IL-2 cytokine release also was evaluated using methods described in Koizumi, et al., 103, 469-475 (1986), which employ trypsinized Jurkat cells rather than PBMNL. Test solutions of the extract were prepared at 100 µg/ml, 10 µg/ml, 1 µg/ml, 0.1 µg/ml and 0.01 µg/ml in 0.1% DMSO.

30

35

Each of the test solutions was incubated overnight, in the presence or absence of co-stimulation by 1 µg/ml calcium ionophore (A23187) and 25 ng/ml PMA (phorbol 12-myristate 13-acetate), with trypsinized Jurkat (human T lymphoid) cells ( $2 \times 10^6$ /ml) suspended with 10% fetal bovine serum in RPMI-1640 at a pH of 7.4 and a temperature of 37°C. The cell suspension was subjected to centrifugation, and the supernatant was evaluated for IL-2 release by use of

- 21 -

an IL-2 immunoassay kit. The results of these assays are set forth in Table 5.

## EXAMPLE 12

5

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on IL-6 Cytokine Release

10 The inhibitory effect of the organic extract obtained in Preparation Example 1 on IL-6 cytokine release was evaluated using methods described in Welker, et al., International Arch of Allergy and Immunology, 109, 110-115 (1996). Test solutions of the extract were prepared at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 15 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$  in 0.1% DMSO.

Each of the test solutions was incubated overnight with LPS (25 ng/ml) and stimulated human peripheral blood mononuclear leukocytes (PBMNL) in growth 20 medium RPMI-1640 at a pH of 7.4 and a temperature of 37°C. The IL-6 cytokine production levels in the conditioned medium were quantitated using a sandwich ELISA kit. The results of these assays are set forth in Table 5.

25

## EXAMPLE 13

30 Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on Interferon- $\gamma$  Cytokine Release

30

The inhibitory effect of the organic extract obtained in Preparation Example 1 on Interferon- $\gamma$  (IFN- $\gamma$ ) cytokine release was evaluated using methods described in the following references: (1) Cohen, 35 et al., Am. J. Clin. Pathol. 105, 589-598 (1996); (2) Henderson, et al., TIPS, 13, 145-151 (1992); (3) Welker, et al., International Arch of Allergy and

- 22 -

Immunology, 109, 110-115 (1996); and (4) Elias, et al., J. Immunol., 138, 3812-3816 (1987).

Test solutions of the extract were prepared at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$  in 0.1% DMSO.

Each of the test solutions was incubated overnight with Concanavalin A (Con-A, 10  $\mu\text{g/ml}$ )-stimulated human peripheral blood mononuclear cells (PBMNCs) in RPMI-1640 growth medium at a pH of 7.4 and a temperature of 37°C. IFN- $\gamma$  cytokine levels in the conditioned medium were then quantitated using a sandwich ELISA kit. The results of these assays are set forth in Table 5.

#### EXAMPLE 14

##### Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on TNF- $\alpha$ Cytokine Release

The inhibitory effect of the organic extract obtained in Preparation Example 1 on TNF- $\alpha$  cytokine release was evaluated using methods described in the following references: (1) Cohen, et al., Am. J. Clin. Pathol. 105, 589-598 (1996); (2) Henderson, et al., TIPS, 13, 145-151 (1992); and (3) Welker, et al., International Arch of Allergy and Immunology, 109, 110-115 (1996). Test solutions of the extract were prepared at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$  in 0.1% DMSO.

Each of the test solutions was incubated overnight with LPS (25 ng/ml) and stimulated human peripheral blood mononuclear cells (PBMNCs) in RPMI-1640 growth medium at a pH of 7.4 and a temperature of 37°C. TNF- $\alpha$  cytokine levels in the conditioned medium were

- 23 -

then quantitated using a sandwich ELISA kit. The results of these assays are set forth in Table 5.

## EXAMPLE 15

5

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on TNF- $\alpha$  Mediated PGE<sub>2</sub> Release

10 The inhibitory effect of the organic extract obtained in Preparation Example 1 on TNF- $\alpha$  mediated PGE<sub>2</sub> release was evaluated using methods described in Lenardo, et al., Cell, 58, 227-229 (1989). Test solutions of the extract were prepared at 100  $\mu$ g/ml, 10  $\mu$ g/ml, 1  $\mu$ g/ml, 0.1  $\mu$ g/ml and 0.01  $\mu$ g/ml in 0.1% DMSO.

15

Each of the test solutions was incubated overnight in the presence or absence of 25 nM Tumor Necrosis Factor- $\alpha$  (TNF- $\alpha$ ), with trypsinized HeLa (human epithelioid cervix carcinoma) S3 cells (2 x 10<sup>6</sup>/ml) suspended with 10% fetal bovine serum in MEM at a pH of 7.3 and a temperature of 37°C. The cell suspension of each well was then transferred to an Eppendorf vial, subjected to centrifugation, and the supernatant was evaluated for released PGE<sub>2</sub> (prostaglandin E<sub>2</sub>) by radioimmunoassay. The results of these assays are set forth in Table 5.

20

25

## EXAMPLE 16

30

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on IL-1 $\alpha$  Mediated PGE<sub>2</sub> Release

35 The inhibitory effect of the organic extract obtained in Preparation Example 1 on IL-1 $\alpha$  mediated PGE<sub>2</sub> release was evaluated using methods described in

- 24 -

Maloff, et al., Clin. Chim. Acta, 180 73-78 (1989).  
Test solutions of the extract were prepared at 100  
 $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  $\mu\text{g/ml}$   
in 0.1% DMSO.

5 Each of the test solutions was incubated overnight,  
in the presence or absence of 1 nM interleukin- $1\alpha$   
(IL- $1\alpha$ ), with trypsinized WI-38 (human diploid  
fibroblast lung) cells ( $10^6/\text{ml}$ ) suspended with 10%  
10 fetal bovine serum in MEM at a pH of 7.3 and a  
temperature of 37°C. The cell suspension of each  
well was then transferred to an Eppendorf vial,  
subjected to centrifugation, and the supernatant was  
evaluated for released PGE $_2$  (prostaglandin E $_2$ ) by  
15 radioimmunoassay. The results of these assays are  
set forth in Table 5.

## EXAMPLE 17

20 Inhibitory Effect Of Organic Extract of *Atractylodes*  
*lancea* Rhizomes on NF $\kappa$ B

The inhibitory effect of the organic extract  
obtained in Preparation Example 1 on NF $\kappa$ B was  
25 evaluated using the methods described in Karttunen,  
et al., Proc. Nat'l. Acad. Sci. USA, 88, 3972-3976  
(1991). Test solutions of the extract were prepared  
at 100  $\mu\text{g/ml}$ , 10  $\mu\text{g/ml}$ , 1  $\mu\text{g/ml}$ , 0.1  $\mu\text{g/ml}$  and 0.01  
 $\mu\text{g/ml}$  in 0.1% DMSO.

30 Jurkat (human T lymphoid) cells transfected with a  
response element-lacZ reporter in which  
transcription of the  $\beta$ -galactosidase gene is  
directed by the binding site for the NF- $\kappa$ B  
35 transcription factor ( $\kappa$ B-Z cells) were used in these  
assays. Each of the test solutions was incubated  
with  $\kappa$ B-Z cells ( $2 \times 10^5$ ) in the presence of 2  $\mu\text{M}$  of

- 25 -

a calcium ionophore (A23187) and 20 ng/ml PMA in RPMI buffer at a pH of 7.4 for 4 hours at 37°C. The cells were then centrifuged, resuspended in buffer and conversion of FDG (fluorescein di- $\beta$ -D-galactopyranoside) to fluorescein by induced  $\beta$ -galactosidase activity was determined after overnight incubation in the dark at 25°C. Fluorescence intensity was measured using Cytofluor (2300) plate reader with excitation at 485 nm and emission at 530 nm. The results of these assays are set forth in Table 5.

## EXAMPLE 18

Inhibitory Effect Of Organic Extract of *Atractylodes lancea* Rhizomes on NF-AT

The inhibitory effect of the organic extract obtained in Preparation Example 1 on NF-AT was evaluated using the methods described in Emmel, et al., Science, 246, 1617-1620 (1989). Test solutions of the extract were prepared at 100  $\mu$ g/ml, 10  $\mu$ g/ml, 1  $\mu$ g/ml, 0.1  $\mu$ g/ml and 0.01  $\mu$ g/ml in 0.1% DMSO.

Jurkat (human T lymphoid) cells transfected with a response element-lacZ reporter in which transcription of the  $\beta$ -galactosidase gene is directed by the binding site for the NFAT-1 transcription factor were used for the assays. Each of the test solutions was incubated with cells ( $2 \times 10^5$ ) in the presence of 2  $\mu$ M of a calcium ionophore (A23187) and 20 ng/ml PMA in RPMI buffer at a pH of 7.4 for 4 hours at 37°C. The cells were then centrifuged and resuspended in buffer, and conversion of FDG (fluorescein di- $\beta$ -D-galactopyranoside) to fluorescein by induced  $\beta$ -galactosidase activity was determined after

- 26 -

overnight incubation in the dark at 25°C. Fluorescence intensity was measured using a Cytofluor (2300) plate reader with excitation at 485 nm and emission at 530 nm. The results of these assays are set forth in Table 5.

Table 5

Proinflammatory Factor	A. lancea extract conc. ( $\mu\text{g/ml}$ )	% Inhibition	IC <sub>50</sub> ( $\mu\text{g/ml}$ )
15-Lipoxygenase	100	50	
Thromboxane Synthetase	100	88	
Cell Adhesion (Fibronectin)	100	116	5.4
Cell Adhesion (Fibronectin)	10.0	65	
Cell Adhesion (Fibronectin)	1.0	13	
Cell Adhesion (Fibronectin)	0.1	15	
Cell Adhesion (Fibronectin)	0.01	10	
Cell Adhesion (VCAM-1)	100	109	9.9
Cell Adhesion (VCAM-1)	10.0	52	
Cell Adhesion (VCAM-1)	1.0	-11	
Cell Adhesion (VCAM-1)	0.1	9	
Cell Adhesion (VCAM-1)	0.01	13	
Cytokine Release (IL-1 $\beta$ )	100	110	7.2
Cytokine Release (IL-1 $\beta$ )	10.0	85	
Cytokine Release (IL-1 $\beta$ )	1.0	-8	
Cytokine Release (IL-1 $\beta$ )	0.1	-3	
Cytokine Release (IL-1 $\beta$ )	0.01	-9	
Cytokine Release (IL-2) PBMNL	100	102	0.4549
Cytokine Release (IL-2) PBMNL	10.0	89	
Cytokine Release (IL-2) PBMNL	1.0	45	
Cytokine Release (IL-2) PBMNL	0.1	37	
Cytokine Release (IL-2) PBMNL	0.01	20	



- 27 -

	Proinflammatory Factor	A. lancea extract conc. ( $\mu\text{g/ml}$ )	% Inhibition	IC <sub>50</sub> ( $\mu\text{g/ml}$ )
5	Cytokine Release (IL-2) Jurkat	100	87	70.6
	Cytokine Release (IL-2) Jurkat	10.0	-13	
	Cytokine Release (IL-2) Jurkat	1.0	-11	
	Cytokine Release (IL-2) Jurkat	0.1	-9	
	Cytokine Release (IL-2) Jurkat	0.01	-9	
10	Cytokine Release (IL-6)	100	100	2.7
	Cytokine Release (IL-6)	10.0	99	
	Cytokine Release (IL-6)	1.0	3	
	Cytokine Release (IL-6)	0.1	-4	
	Cytokine Release (IL-6)	0.01	-2	
15	Cytokine Release (Interferon- $\gamma$ )	100	99	5.7
	Cytokine Release (Interferon- $\gamma$ )	10.0	68	
	Cytokine Release (Interferon- $\gamma$ )	1.0	9	
	Cytokine Release (Interferon- $\gamma$ )	0.1	-2	
	Cytokine Release (Interferon- $\gamma$ )	0.01	4	
20	Cytokine Release (TNF- $\alpha$ )	100	107	9.3
	Cytokine Release (TNF- $\alpha$ )	10.0	59	
	Cytokine Release (TNF- $\alpha$ )	1.0	-18	
	Cytokine Release (TNF- $\alpha$ )	0.1	-13	
	Cytokine Release (TNF- $\alpha$ )	0.01	-2	
25	TNF- $\alpha$ Mediated PGE <sub>2</sub> Release	100	90	
	TNF- $\alpha$ Mediated PGE <sub>2</sub> Release	10.0	20	
	TNF- $\alpha$ Mediated PGE <sub>2</sub> Release	1.0	-5	
	TNF- $\alpha$ Mediated PGE <sub>2</sub> Release	0.1	-2	
	TNF- $\alpha$ Mediated PGE <sub>2</sub> Release	0.01	0	
30	IL-1 $\alpha$ Mediated PGE <sub>2</sub> Release	100	110	
	IL-1 $\alpha$ Mediated PGE <sub>2</sub> Release	10.0	44	
	IL-1 $\alpha$ Mediated PGE <sub>2</sub> Release	1.0	20	
	IL-1 $\alpha$ Mediated PGE <sub>2</sub> Release	0.1	8	
	IL-1 $\alpha$ Mediated PGE <sub>2</sub> Release	0.01	-1	

- 28 -

Proinflammatory Factor	A. lancea extract conc. ( $\mu\text{g/ml}$ )	% Inhibition	IC <sub>50</sub> ( $\mu\text{g/ml}$ )
NF $\kappa$ B transcription	100	96	7
NF $\kappa$ B transcription	10.0	59	
NF $\kappa$ B transcription	1.0	11	
NF $\kappa$ B transcription	0.1	7	
NF $\kappa$ B transcription	0.01	-13	
NF-AT transcription	100	96	8.8
NF-AT transcription	10.0	54	
NF-AT transcription	1.0	7	
NF-AT transcription	0.1	-12	
NF-AT transcription	0.01	-10	

## EXAMPLE 19

A mammal having or at risk for developing a condition which is benefited by inhibition of the activity of COX-2 or other proinflammatory factors may be treated with a therapeutically or prophylactically effective amount of the *Atractylodes lancea* extract of this invention.

The therapeutically effective or prophylactically effective amount of *Atractylodes lancea* extract may be administered to the mammal by any of the known routes for administering a pharmaceutical agent to a patient, including parenterally, orally and rectally. The *Atractylodes lancea* extract may be administered using a dosing regimen which is effective to inhibit the activity of COX-2 or other proinflammatory factors in the mammal for a desired period of time. The proper dosage amount may be determined by routine experimentation using methods known in the art.

- 29 -

Other variations and modifications of this invention will be obvious to those skilled in the art. This invention is not limited, except as set forth in the claims.

- 30 -

## WHAT IS CLAIMED IS:

1. A method for inhibiting the activity of a proinflammatory factor in a mammal, said method comprising the step of administering to the mammal an organic extract of *Atractylodes lancea*.
2. A method according to claim 1, wherein said proinflammatory factor is selected from the group consisting of cyclooxygenase-2, 15-lipoxygenase, thromboxane synthetase, inflammatory cell adhesion to fibronectin, inflammatory cell adhesion to VCAM-1, IL-1 $\beta$  cytokine release, IL-2 cytokine release, IL-6 cytokine release, Interferon- $\gamma$  cytokine release, TNF- $\alpha$  cytokine release, TNF- $\alpha$  mediated PGE<sub>2</sub> release, IL-1 $\alpha$  mediated PGE<sub>2</sub> release, NF-AT transcription of proinflammatory genes, and NF- $\kappa$ B transcription of proinflammatory genes.
3. A method according to claim 1, wherein said proinflammatory factor is cyclooxygenase-2.
4. A method according to claim 3, wherein the inhibitory effect of said extract on cyclooxygenase-2 is greater than or equal to about 2 times greater than the inhibitory effect of said extract on cyclooxygenase-1.
5. A method according to claim 3, wherein the inhibitory effect of said extract on cyclooxygenase-2 is greater than or equal to about 10 times greater than the inhibitory effect of said extract on cyclooxygenase-1.
6. A method according to any of claims 1-5, wherein said organic extract is a purified compound obtained by a process comprising the steps of:

- 31 -

(i) exposing rhizomes of *Atractylodes lancea* to an organic solvent under conditions appropriate to remove a proinflammatory factor inhibitory extract from said rhizomes; and

(ii) isolating said proinflammatory factor inhibitory extract.

7. A method according to claim 6, wherein step (i) comprising mixing said rhizomes with said solvent and stirring the resultant mixture at a temperature between about 25°C and the boiling point of said solvent for at least one minute.

8. A method according to claim 7, wherein said solvent is an organic solvent.

9. A method according to claim 8, wherein said organic solvent is selected from the group consisting of hydrocarbon solvents, ethers, chlorinated solvents, acetone, ethyl acetate, butanol, ethanol, methanol, isopropyl alcohol and mixtures thereof.

10. A method according to claim 9, wherein said nonpolar organic solvent is dichloromethane.

11. A method according to claim 9, wherein step (ii) comprises separating said solvent from said organic extract by evaporating said solvent.

12. A method according to claim 6, comprising administering a therapeutically effective or prophylactically effective amount of said purified compound to a mammal which has or is at risk for developing a condition which is benefited by the inhibition of the activity of a proinflammatory factor.

- 32 -

13. A method according to claim 12, wherein said condition is a cyclooxygenase-2 mediated condition.

14. A method according to claim 13, wherein said cyclooxygenase-2 mediated condition is general inflammation.

15. A method according to claim 13, wherein said cyclooxygenase-2 mediated condition is arthritis.

16. A method according to claim 13, wherein said cyclooxygenase-2 mediated condition is pain.

17. A method according to claim 13, wherein said cyclooxygenase-2 mediated condition is cancer.

18. A method according to claim 12, wherein said purified compound is administered as a pharmaceutical composition comprising a pharmaceutically acceptable excipient.

19. A method according to claim 12, wherein said purified compound is administered as a nutritional composition comprising a nutritionally acceptable excipient.

20. A method according to claim 12, wherein said purified compound is administered as a food composition.

21. A method according to claim 12, wherein said purified compound is administered as a food ingredient composition.

1/7






FIG. 1

2 / 7



A. LANCEA



SUBSTITUTE SHEET (RULE 26)






FIG. 4




FIG. 5





FIG. 6

SUBSTITUTE SHEET (RULE 26)





FIG. 7

SUBSTITUTE SHEET (RULE 26)

6/7




FIG. 8

SUBSTITUTE SHEET (RULE 26)




FIG. 9

SUBSTITUTE SHEET (RULE 26)

# INTERNATIONAL SEARCH REPORT

In ternational Application No  
PCT/US 00/07085

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61K35/78 A61P29/02

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 613 591 A (ABURADA MASAKI ET AL) 23 September 1986 (1986-09-23) column 2, line 38 - line 50; claims ---	1-21
X	DATABASE WPI Section Ch, Week 199237 Derwent Publications Ltd., London, GB; Class B03, AN 1992-303541 XP002142228 & JP 04 208222 A (TSUMURA & CO), 29 July 1992 (1992-07-29) abstract --- -/--	1-21

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

10 July 2000

Date of mailing of the international search report

25/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Seegert, K

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/07085

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 199208 Derwent Publications Ltd., London, GB; Class B04, AN 1992-060692 XP002142229 & JP 04 005237 A (NONOKAWA SHOJI YG), 9 January 1992 (1992-01-09) abstract ---	1-21
Y	RESCH M. ET AL: "5-Lipoxygenase and cyclooxygenase-1 inhibitory active compounds from Atractylodes lancea" J. NAT. PROD., vol. 61, 1998, pages 347-250, XP000916172 cited in the application abstract ---	1-21
Y	DATABASE MEDLINE 'Online! AN: 94178130, 1993 XP002142227 abstract & CHANG M L ET AL: "The influence of Chinese traditional medicine on the production and activity of interleukin 1 (IL-1)" JOURNAL OF MICROBIOLOGY AND IMMUNOLOGY , vol. 26, no. 1, 1993, pages 15-24, -----	1-21

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/07085

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4613591 A	23-09-1986	JP 1633501 C	20-01-1992
		JP 2056329 B	29-11-1990
		JP 59036619 A	28-02-1984
		AT 383740 B	10-08-1987
		AT 299783 A	15-01-1987
		BE 897403 A	14-11-1983
		CH 659948 A	13-03-1987
		DE 3328263 A	23-02-1984
		FR 2531864 A	24-02-1984
		GB 2126085 A, B	21-03-1984
		HU 187103 B	28-11-1985
		IT 1169771 B	03-06-1987
		NL 8302391 A	16-03-1984
JP 4208222 A	29-07-1992	NONE	
JP 4005237 A	09-01-1992	NONE	